arXiv:quant-ph/0105034v4 22 Nov 2001

Four-photon Entanglement as Stochastic-signal Correlation

A. F. Kracklauer
kracklau@fossi.uni-weimar.de

A fully classical model of a recent experiment exhibiting what is interpreted as teleportation and four-photon entanglement is described. It is shown that the reason that a classical model is possible, contrary to the current belief, results ultimately from a misguided modification by Bohm of the EPR *Gedanken* experiment. Finally, teleportation is reinterpreted as the passive filtration of correlated but stochastic events in stead of the active transfer of either material or information.

PACS numbers: 03.65.Bz, 03.67.-a, 41.10.Hv, 42.50.Ar

In a recent letter J.-W. Pan et al. described a demonstration of the experimental observation "of pure four-photon GHZ entanglement produced by parametric down-conversion and a projective measurement."[1] They add that this experiment "demonstrates teleportation with very high purity" and that "the high visibility not only enables various novel tests of quantum nonlocality, [but] it also opens the possibility to experimentally investigate quantum computation and communications schemes with linear optics." This demonstration was achieved using a novel and ingeniously simple (in concept, not necessarily in concrete realization) setup employing two sources of entangled photons which feed the two faces of a polarizing beam splitter (PBS). The authors further claim that this setup gives results that finally preclude all doubt that nonlocal effects ensue from quantum entanglement.

It is the purpose of this letter to contest this last claim. This will be achieved by proposing a classical model of their experiment, without 'nonlocality,' which fully duplicates its results.

First, we briefly review the experimental setup. (See Fig. 1) Two independent entangled photon pairs are created by down-conversion in a crystal pumped by a pulsed laser. The laser pulse passes through the crystal creating one pair (A), then is reflected off a movable mirror to repass through the crystal in the opposite direction creating a second pair (B). One photon from each pair is fed directly through polarizers to photodetectors (photons 1 and 4). The other photons (2 and 3) are directed to opposite faces of a PBS, (i.e., a beam splitter which reflects vertically and transmits horizontally polarized photons) after which the exiting photons are sent through variable polarizers into photodetectors. The path lengths for photons 2 and 3 are adjusted so as to compensate for the time delay in the creation of the pairs. By moving the mirror, the compensation can be negated to permit observing the disappearance of interference caused by lack of simultaneous "cross-talk" between channels 2 and 3.

The reported observations are the following: Of all the 16 possible polarizer settings regimes for which $\theta_n = 0$ or $\pi/2$ w.r.t. the horizontal axis of the PBS, only $\{0,\ \pi/2,\ \pi/2,\ 0\}$ and $\{\pi/2,\ 0,\ 0,\ \pi/2\}$ yield a (substantial) four-fold coincidence count: C; the regime $\{\pi/4,\ \pi/4,\ \pi/4,\ \pi/4\}$ occurs with an intensity $C/4.2$ and the regime $\{\pi/4,\ \pi/4,\ \pi/4,\ -\pi/4\}$ with zero intensity. Further, both of the latter regimes yield an intensity of $C/8$ when photons 2 and 3 do not overlap.



FIG. 1: Schematic of the experimental setup for the measurement of four-photon GHZ correlations. A pulse of laser light passes a nonlinear crystal twice to produce two entangled photon pairs via parametric down conversion. Coincidences between all four detectors are used to study the nature of entanglement.

The model proposed herein, like all others, is based on certain assumptions, which, being quite different from those couched in the notation and vocabulary of QM, must be delineated explicitly. They consist of the following:

1. Electromagnetic radiation is comprised of continuous waves as described by Maxwell's equations.
2. All detectors in the optical region of the electromagnetic spectrum exploit the photoelectric effect. They convert continuous radiation to an electron current. Electrons are discrete objects; their generation in a photodetector effectively digitizes the signal associated with incoming radiation, thereby evoking the impression that the radiation was itself somehow digitized into units (photons). The last inference is unwarranted; we can not know what form the incoming energy actually had, we are restricted to inferring its nature from the electron

current. It is known empirically that photoelectrons are ejected from a photodetector randomly but in proportion to the energy density, E^2, of the incoming signal. For the model, we take the conversion efficiency to be 100%.

3. The nonlinear crystal generating signals by parametric down-conversion is taken to be a "black box" which, by virtue of its structure emits randomly pairs of signal pulses confined to the pure vertical and horizontal polarization modes. The pairs are anticorrelated with respect to polarization and each member of a pair is directed (in the well known manner) into two intersecting cones; all horizontal emissions are into one cone, all vertical into the other. The two variants of pairs are taken to be equally likely. "Entangled" radiation samples extracted from the two points where the cones intersect, therefore, constitute a mirror image pair of random sequences of individual electromagnetic pulses of both pure modes.
4. The intensity of four-fold coincidence detections among four photodetectors is calculated using non quantum coherence theory.[2] Coincident count probabilities, for a system with N, (herein $N = 4$), monitored exit ports are proportional to the single time, multiple location second order (in intensity) cross correlation, i.e.:

$$P(\theta_1,\, \theta_2, ..\theta_N) = \frac{< \prod_{n=1}^{N} E^*(r_n,\, \theta_n) \prod_{n=N}^{1} E(r_n,\, \theta_n) >}{\prod_{n=1}^{N} < E^*(r_n) E_n(r_n) >}. \tag{1}$$

For this model the denominator consists of constants of factors of the form $(\cos^2(a) + \sin^2(a))$ so that it equals 1.

These assumptions, all fully compatible with classical physics, account for all of the reported observations. Eq. (1) was implemented numerically as follows: The centers are assumed to emit double pulses in opposed directions which are anticorrelated and confined to the vertical and horizontal polarization modes; i.e. their polarization vectors are:

$$\begin{aligned} A_1 &= (cos(n\frac{\pi}{2}),\, sin(n\frac{\pi}{2})), & (2)\\ A_2 &= (sin(n\frac{\pi}{2}),\, -cos(n\frac{\pi}{2})), & (3)\\ B_1 &= (cos(m\frac{\pi}{2}),\, sin(m\frac{\pi}{2})), & (4)\\ B_2 &= (sin(m\frac{\pi}{2}),\, -cos(m\frac{\pi}{2})), & (5) \end{aligned}$$

where n and m take the values 0 and 1 with a flat random distribution. The polarizing beam splitter (PBS) is modeled using the transition matrix for a polarizer,

$$P(\theta) = \begin{bmatrix} \cos^2(\theta) & \cos(\theta)\sin(\theta) \\ \sin(\theta)\cos(\theta) & \sin^2(\theta) \end{bmatrix}, \tag{6}$$

where $\theta = \pi/2$ accounts for a reflection, and $\theta = 0$ a transmission. Thus, the field impinging on each of the four detectors is:

$$\begin{aligned} E_1 &= P(\theta_1)A_1, & (7)\\ E_2 &= P(\theta_2)(P(0)B_2 - P(\pi/2)A_2), & (8)\\ E_3 &= P(\theta_3)(P(0)A_2 - P(\pi/2)B_2), & (9)\\ E_4 &= P(\theta_4)B_1. & (10) \end{aligned}$$

Putting Eqs. (7-10) into Eq. (1) and carrying out the indicated average by summing on n and m and dividing by 4 as there are four combinations, one obtains the expression for the fourth order (in fields) coincidence correlation as a function of the polarizer settings. The result is cumbersome but easily evaluated numerically for various regimes, i.e., various selections of θ_n. Ideal results compatible to those observed are obtained. Slight deviations in the observed data appear to be due to noise.



FIG. 2: The upper curve shows the effect on the intensity of four-fold coincidences of skewing (rotating) all polarizers through a given angle starting from the state $\{\pi/2, 0, 0, \pi/2\}$. The lower curve shows the same effect when one of the polarizers is rotated in the opposite direction. The middle curve shows the effect of either of these skewing schemes when the timing is such that the crossover signals do not arrive simultaneously with the reflected signals. Note that the separation of the curves and values at $\pi/4$ coincide with the observations reported in Ref. [1]. This diagram differs from Fig. 4 in Ref. [1] in that it shows the split of these regimes as a function of polarizer skew for fixed delay rather than as a function of delay for fixed skew.

To model regimes for which the pulse pairs were generated with a time difference such that the cross-over signals in beams 2 and 3 could not interfere, the sum on the radiation contributions from individual centers is squared *before* averaging. This procedure recognizes the fact that the electric fields from distinct sources (A and B, in this case) must be added *after* squaring because they do not interfere but still deposit energy into a photodetector. Changing the order of 'squaring' and 'summing' for this setup affects only the crossover signals as all others are in orthogonal polarization modes in any case. All this leads, again, to results exactly mimicking the effects reported. Additionally, the relative count intensity in other regimes can be computed easily; for an example, see Fig. 2.

Note that on Fig. 2 the curve for the 'state' $< \pi/4, \pi/4, \pi/4, \pi/4 >$ splits or separates from the 'state' $< \pi/4, \pi/4, \pi/4, -\pi/4 >$ at $\theta = \pi/4$. Pan et al. interpret this splitting to indicate: "that the state of, say, photon 2 was teleported to photon 4..." and "...fully demonstrates the nonlocal feature of quantum teleportation." The mere fact that this splitting is faithfully modeled classically, using, e.g., coherent vice Fock states, casts strong doubt on such inferences.

There is nothing essentially quantum mechanical in this model. It does not make use of any property peculiar to QM. In particular it does not in any way assume properties of "photons" and its results do not depend, therefore, on assumptions regarding detector efficiency (effectively taken to be 100%). It constitutes a fully faithful classical technique for modeling experiments carried out in a portion of the spectrum admitting macroscopic devices and detailed time tracking of electromagnetic fields, thereby evading the peculiarities of photodetectors. (Note that EPR-B correlations have been so observed.[3]) The same type of model for EPR-B experiments yields exactly the same expressions as those given by QM which violate the relevant Bell Inequality.

In view of the conventional wisdom that it is impossible to comprehend phenomena involving *quantum* entanglement using non quantum physics [4], it is natural to ask how it is possible then that a classical model can be suitable. We believe that the genesis of this misunderstanding is to be found with D. Bohm.[5] The over-arcing issue ultimately motivating the experiment described in Ref. [1], and further analyzed herein is the Einstein, Podolsky and Rosen [EPR] argument of 1935 to the effect that QM is incomplete.[6] In that paper EPR argued that the position and momentum of two entangled particles, created auspiciously, could be specified exactly in spite of Heisenberg Uncertainty, by exploiting symmetry. With this argument, they hoped to show that Heisenberg Uncertainty was not something fundamentally new, but just ignorance. This being the case, they argued, there should exist a deeper theory, involving heretofore "hidden variables" that would more precisely describe nature. The point of their whole consideration involved the ultimate nature of Heisenberg Uncertainty.

Much later Bohm modified EPR's original *Gedanken* Experiment. He, for reasons of simplicity, transfered the EPR argument from phase space, (x, p) to another arena, one involving spin.[5] This too turned out to be experimentally unrealizable, but the algebraically isomorphic arena involving polarized light is entirely practical. Thus, nowadays, the EPR issue is discussed, analyzed and explored in terms of polarization phenomena. However, note that because of Heisenberg Uncertainty, (x, p) do *not* commute, i.e., $[\hat{x}, \hat{p}] = i\hbar$, whereas the basis operators of polarization space $\hat{E}_h$, $\hat{E}_p$, where $\hat{E}_x$ represents an electric field in the x direction, *do* commute. That is, there is no Heisenberg Uncertainty among polarization modes. This is a fact substantiated by QM itself; creation and annihilation operators for different modes of polarization, commute. Bohm's transfer of venue moved the issue from one in which there is Heisenberg Uncertainty into one in which there is not! Experiments in the polarization arena can not, therefore, address the issue introduced by EPR; they, by logical necessity, leave it unexamined. It is simply impossible to investigate Heisenberg Uncertainty where there is none.

Because the basis operators of polarization space (a.k.a. "qubit space") commute, all polarization phenomena ultimately must be describable with non quantum principles. It is for this reason, that a classical model can explain this experiment. Alternately, this conclusion follows forthwith, albeit with mostly only formalistic authority, from the Optical Equivalence Theorem.[7]

Bohm did not justify carefully his change of venue; he simply declared spin operators to be equivalent and sallied forth. This has been accepted, apparently, on the grounds that, like the basis operators of phase space, (x, p) , the spin operators (σ_x, σ_y) also do not commute. While this is indeed true, the reason is not the same. Spin operators with discrete eigenvalues pertain to the direction in space defined by a magnetic field. In directions transverse to the magnetic field, the expectation values are not discrete but oscillate out of phase. In the end, the reason spin operators do not commute would be that it is impossible to have more than one direction for a magnetic field or precession axis at a time. Non commutation, then, is not a manifestation of Heisenberg Uncertainty in this case, but of geometry, i.e., of the nature of an axis of rotation. Indeed, angular momentum operators do not commute in classical mechanics. Similar remarks pertain to the privileged direction defined by the "**k**" vector of an EM wave with respect to the transverse, i.e., polarization directions.

Entanglement is often cited as the core of QM, an idea for which perhaps Schrödinger was the originator.[8] However, if entanglement is defined in terms of the non factorability of a wave function, as it most often is, then it must be attributable to a correlation between subsystems.Obviously, if a wave function factors into terms each pertaining to a separate subsystem, $\psi_1(r_1)\psi_2(r_2)$, then the probability density computed from this product, $\psi_1^*\psi_2^*(r_1, r_2)\psi_2\psi_1(r_1, r_2) = (\psi_1^*(r_1)\psi_1(r_1))(\psi_2^*(r_2)\psi_2(r_2))$, also factors, in which case it pertains to *statistically independent,* i.e., uncorrelated subsystems. If the wave function does not factor, then the probability density also will not factor and, clearly, the subsystems are simply not uncorrelated. Correlation need not imply nonlocality; hereditary correlation satisfies all requirements posed by the physical situation. In experiments involving EPR style setups, including that described herein, correlation can be vested in the 'pho-

```
> restart:  a:=1:  b:=1:  c:=0:  d:=0:  e:=1:
> with(linalg):
> Pro:=z->matrix([[cos(z)^2,cos(z)*sin(z)],[sin(z)*cos(z),sin(z)^2]]):
> A1:=vector([cos(n*Pi/2),sin(n*Pi/2)]):
> A2:=vector([sin(n*Pi/2),-cos(n*Pi/2)]):
> B1:=vector([cos(m*Pi/2),sin(m*Pi/2)]):
> B2:=vector([sin(m*Pi/2),-cos(m*Pi/2)]):
> E1:=multiply(Pro(z1),A1):
> E2:=multiply(Pro(z2),-multiply(Pro(Pi/2),A2)+multiply(Pro(0),B2)):
> E3:=multiply(Pro(z3),-multiply(Pro(Pi/2),B2)+multiply(Pro(0),A2)):
> E4:=multiply(Pro(z4),B1):
> EE:=sum(sum(sum(sum((E1[i]*E2[j]*E3[k]*E4[l]),i=1..2),j=1..2),k=1..2),l=1..2):
> TrJ:=seq(sum(sum(op(v,EE)^(2-e),n=0..1),m=0..1)^(1+e),v=1..16):
> z1:=c*Pi/2+a*t:  z2:=d*Pi/2+b*t:  z3:=d*Pi/2+t:  z4:=c*Pi/2+t:
> In:=sum(TrJ[q],q=1..16):
> plot(In(t),t=0..Pi):
```

FIG. 3: A Maple implementation of Eq. (1). The polarizer regimes are encoded by setting the values of a, b, c and d to ± 1, 0. 't' is the skew angle. Setting the value of e to 1, adds the EM field contribution from the various centers before 'squaring;' as physics, this accounts for "crossover" signals that arrive simultaneously with the reflected signals and interfere. Setting e to zero, on the other hand, prevents interference of these signals. (This works with this model because of the peculiarities of down-conversion and a PBS; care must be taken modeling other setups to (in-)exclude appropriate signals before squaring.)

tons,' or in classical terms: the signals, at their origin and simply carried along thereafter.

The actual problem with entanglement arises elsewhere — with *particle* beams. A wave function describing a particle beam can not be considered simply as pertaining to a physical ensemble, because particles, one by one, are diffracted at slits, for example; they suffer, seemingly, 'entanglement' between Gibbsian ensemble members. Rationalizing this phenomenon requires other arguments[9]; but, the desideratum of uniformity of principles, has lead to the mandate that radiation, too, be considered ontologically ambiguous until the moment of measurement, even though there is no need to do so. Its apparent digitization can be seen simply as a manifestation of the nature of photoelectron detectors.

Of course, the tactic employed herein works for all phenomena for which there is no Heisenberg Uncertainty, e.g., simple EPR correlations.[10] It offers a decidedly less mystical interpretation of many phenomena typically ascribed to QM. Teleportation, for example, admits a passive interpretation involving no "portation" of any nature. In the above model, so-called teleported states (1 and 4) are those which, although from separate random sources, eventually match up, and this is signaled by an appropriate coincidence between each's partner (2 and 3). Such an explanation is decidedly less enchanting than that conveyed by the term "teleportation," but hugely more respectful of principles desirable for a rational explanation of the natural world.

[1] J.-W. Pan, M. Daniell, S. Gasparoni, G. Weihs and A. Zeilinger, *Phys. Rev. Lett.* **86**, 4435, (2001).

[2] L. Mandel and E. Wolf, *Optical Coherence and Quantum Optics,* (Cambridge University, Cambridge, 1995).

[3] N. V. Evdokimov, D. N. Klyshko, V. P. Komolov and V. A. Yarochkin, *Physics - Uspekhi* **39,** 83 (1996).

[4] D. Bouwmeester, et al. *The Physics of Quantum Information,* (Springer, Berlin, 2000).

[5] D. Bohm, *Quantum Theory,* (Prentice-Hall, New York, 1951).

[6] A. Einstein, B. Podolsky and N. Rosen, *Phys. Rev.* **47**, 777 (1935).

[7] J. R. Klauder and E. C. G. Sudarshan, *Fundamentals of Quantum Optics*, (Benjamin, New York, 1968), p. 192.

[8] E. Schrödinger, *Naturwissenschaften,* **23,** 807, 823, 844 (1935).

[9] A. F. Kracklauer, *Found. Phys. Lett.* **12** (5) 441 (1999).

[10] A. F. Kracklauer, *Ann. Fond. L. deBroglie* **20** (2) 193, (2000); quant-ph/0108057.